

Prescribing information and adverse event reporting information for EYLEA® (aflibercept) can be accessed [here](#) or via the QR code located on the last page of this document.



STREAMLINE YOUR CLINIC WITH EYLEA® (aflibercept) 8 mg



EYLEA 8 mg DOSING REGIMENS IN nAMD AND DMO

EYLEA 8 mg is licensed up to 6 months intervals,* allowing you to extend stable patients out to as few as 2 injections per year.¹⁻⁵

***Following 3 monthly loading doses, intervals can be extended to Q16, dependent on visual and/or anatomic outcomes, and subsequently to Q24 (e.g. with a T&E regimen) if visual and/or anatomic outcomes are stable. Consult the SmPC for full posology.¹**

EYLEA 114.3 mg/mL is indicated in adults for the treatment of neovascular (wet) age-related macular degeneration (nAMD) and visual impairment due to diabetic macular oedema (DMO).¹

For UK healthcare professionals only.

DMO, diabetic macular oedema. nAMD, neovascular age-related macular degeneration. Q16, every 16 weeks. Q24, every 24 weeks. SmPC, Summary of Product Characteristics. T&E, treat and extend. VEGF, vascular endothelial growth factor.

References

1. EYLEA® 114.3 mg/mL Summary of Product Characteristics. **2.** Lanzetta P, et al. Intravitreal aflibercept 8 mg in neovascular age-related macular degeneration (PULSAR): 48-week results from a randomised, double-masked, non-inferiority, Phase 3 trial. *Lancet*. 2024;403(10432):1141-1152. **3.** Brown DM, et al. Intravitreal aflibercept 8 mg in diabetic macular oedema (PHOTON): 48-week results from a randomised, double-masked, non-inferiority, Phase 2/3 trial. *Lancet*. 2024;403(10432):1153-1163. **4.** Lanzetta P. Intravitreal aflibercept 8 mg injection in patients with neovascular age-related macular degeneration: 60-week and 96-week results from the Phase 3 PULSAR trial. *EURETINA*. 5-8 October 2023. Amsterdam, The Netherlands. Oral presentation. **5.** Wykoff CC. Aflibercept 8 mg for diabetic macular edema: 96-week results from the Phase 2/3 PHOTON trial. *EURETINA*. 5-8 October 2023. Amsterdam, The Netherlands. Oral presentation.

TREAT YOUR ELIGIBLE PATIENTS WITH FLEXIBILITY

Extend up to 6-month intervals* and experience flexibility with optional reloading for patients switching from other anti-VEGF agents[†]


For patients initiating treatment, EYLEA 8 mg can be used as part of a T&E regimen to lengthen intervals[‡]

Tx-naïve

Switch

Treatment-naïve patients:

Weeks



After reaching Q16, intervals can be further extended up to Q24

Allowing you to extend stable patients out to as few as 2 injections per year^{†‡}

The frequency of monitoring visits should be based on the patient's status and at the physician's discretion.¹

Dosing diagram for illustrative purposes only.

*Following 3 monthly loading doses, intervals can be extended to Q16, dependent on visual and/or anatomic outcomes, and subsequently to 6 months (e.g. with a T&E regimen) if visual and/or anatomic outcomes are stable.¹

[†]Stable patients switching to EYLEA 8 mg do not require loading doses and can maintain or extend previous treatment intervals after the first injection of EYLEA. Consult the SmPC for full posology.¹

[‡]For patients who have previously been treated with EYLEA 2 mg or other anti-VEGF medicinal products and are switching to EYLEA 8 mg, the treatment regimen can differ from that used for treatment-naïve patients. Treatment intervals should be determined based on visual and/or anatomic outcomes: in patients with stable visual and anatomic outcomes, previous treatment intervals can be maintained or extended after the first injection of EYLEA 114.3 mg/mL, such as with a T&E dosing regimen; in patients with suboptimal visual and/or anatomic outcomes, treatment with EYLEA 8 mg may begin with 1 injection per month for up to 3 consecutive doses followed by adjustment of injection intervals, such as with a T&E dosing regimen.¹

Q16, every 16 weeks; Q24, every 24 weeks; SmPC, Summary of Product Characteristics; T&E, treat and extend; Tx, treatment; VEGF, vascular endothelial growth factor.

Reference: 1. EYLEA® 114.3 mg/mL Summary of Product Characteristics.

TREAT YOUR ELIGIBLE PATIENTS WITH FLEXIBILITY

Extend up to 6-month intervals* and experience flexibility with optional reloading for patients switching from other anti-VEGF agents[†]

Patients who have previously been treated with EYLEA 2 mg or other anti-VEGF medicinal products can maintain or extend their current interval when switching to EYLEA 8 mg, with or without reloading[‡]

Tx-naïve

Switch

Stable visual and anatomic outcomes
(No loading doses required):

Discontinue previous anti-VEGF

Weeks

0

Suboptimal visual and/or anatomic outcomes
(Optional 3 loading doses):

Discontinue previous anti-VEGF

Weeks

0

4

8

Up to 3 consecutive loading doses

Maintain or extend previous treatment interval
(e.g. using T&E up to Q24 intervals)

The frequency of monitoring visits should be based on the patient's status and at the physician's discretion.¹

Dosing diagram for illustrative purposes only.

*Following 3 monthly loading doses, intervals can be extended to Q16, dependent on visual and/or anatomic outcomes, and subsequently to 6 months (e.g. with a T&E regimen) if visual and/or anatomic outcomes are stable.¹

[†]Stable patients switching to EYLEA 8 mg do not require loading doses and can maintain or extend previous treatment intervals after the first injection of EYLEA. Consult the SmPC for full posology.¹

[‡]For patients who have previously been treated with EYLEA 2 mg or other anti-VEGF medicinal products and are switching to EYLEA 8 mg, the treatment regimen can differ from that used for treatment-naïve patients. Treatment intervals should be determined based on visual and/or anatomic outcomes: in patients with stable visual and anatomic outcomes, previous treatment intervals can be maintained or extended after the first injection of EYLEA 114.3 mg/mL, such as with a T&E dosing regimen; in patients with suboptimal visual and/or anatomic outcomes, treatment with EYLEA 8 mg may begin with 1 injection per month for up to 3 consecutive doses followed by adjustment of injection intervals, such as with a T&E dosing regimen.¹

Q16, every 16 weeks; Q24, every 24 weeks; SmPC, Summary of Product Characteristics; T&E, treat and extend; Tx, treatment; VEGF, vascular endothelial growth factor.

Reference: 1. EYLEA® 114.3 mg/mL Summary of Product Characteristics.

Prescribing information and adverse event reporting information for EYLEA® (aflibercept) is available via the QR code on the right.

Either click [here](#) or scan the QR code for prescribing information and adverse event reporting information.

For direct access to this prescribing information, please ensure your device's browser settings have automatic PDF download enabled.

