



Welcome to precision, delivered with OcuClick®

How to use OcuClick, the EYLEA® 8 mg (aflibercept 114.3 mg/mL) solution for injection in pre-filled syringe¹

1. Prepare

When ready to administer EYLEA 8 mg, open the carton and remove the blister pack. Carefully peel open the blister pack to ensure the sterility of its contents.

Keep the syringe in the sterile tray until you are ready to attach the injection needle. Use aseptic technique to carry out steps 2-9.


2. Remove syringe from pack

Remove the syringe from the sterilised blister pack.

3. Inspect syringe and solution for injection

Do not use the pre-filled syringe if particulates, cloudiness, or discolouration are visible.


Do not use if any part of the pre-filled syringe is damaged or loose, or if the syringe cap is detached from the Luer-lock.



4. Snap off syringe cap

Snap off (do not twist off) the syringe cap by holding the syringe in one hand and the syringe cap with the thumb and forefinger of the other hand.


Do not pull back on the plunger rod.



5. Attach 30G needle


Firmly twist the 30 G x 1/2 inch injection needle onto the Luer-lock syringe tip.

Use of a smaller size needle (higher gauge) than the recommended 30 G x 1/2 inch injection needle is not advised due to risk of increased injection force.



6. Dislodge air bubbles

Holding the syringe with the needle pointing up, check the solution for bubbles. If there are bubbles, gently tap the syringe with your finger until the bubbles rise to the top.




7. Expel air and excess volume to prime

The syringe does not have a dose line because it is designed to set the dose mechanically.

Priming and setting the dose must be done using the following steps.

To eliminate all bubbles and to expel excess drug, slowly depress the plunger rod until it stops, i.e. when the guide on the plunger rod reaches the finger grip.




8. Set to dose

Turn the end of the plunger rod 90 degrees clockwise or counterclockwise until the guide of the plunger rod aligns with the slot. You may hear or feel a **click**.

The dose of EYLEA 8 mg is now set and ready for injection. **Do not push the plunger rod before insertion into the eye.**

When ready to administer EYLEA 8 mg, remove the plastic needle shield.



9. Administer the injection

Insert the needle into the ocular injection site.

Inject the solution by pushing the plunger rod until it stops, i.e. until the guide is completely within the slot. Do not apply additional pressure once the guide is within the slot. It is normal to see a small amount of residual solution left in the syringe after the injection.

The pre-filled syringe is for single dose administration and single use only. After injection, discard the used syringe into a sharps container.

EYLEA 114.3 mg/mL is indicated in adults for the treatment of neovascular (wet) age-related macular degeneration (nAMD) and visual impairment due to diabetic macular oedema (DMO).¹

For UK Healthcare professionals only.

This promotional material was organised and fully funded by Bayer.

1. EYLEA® 114.3 mg/mL Summary of Product Characteristics.

Prescribing information and adverse event reporting information for EYLEA® (aflibercept) is available via the QR code on the right.

Either click [here](#) or scan the QR code for prescribing information and adverse event reporting information.

For direct access to this prescribing information, please ensure your device's browser settings have automatic PDF download enabled.

